

When Do Children and Teens Need Vaccinations?

Age	HepB Hepatitis B	DTaP/Tdap Diphtheria, tetanus, pertussis	Hib <i>Haemophilus influenzae</i> type b	Polio	PCV Pneumococcal conjugate	RV Rotavirus	MMR Measles, mumps, rubella	Varicella Chickenpox	HepA Hepatitis A	HPV Human papillo- mavirus	MCV4 Meningococcal conjugate	Influenza	
Birth	✓												
2 months	✓ (1–2 mos)	✓	✓	✓	✓	✓							
4 months	✓ ¹	✓	✓	✓	✓	✓							
6 months		✓	✓ ²		✓	✓ ²							
12 months	✓ (6–18 mos)	✓ ⁴ (15–18 mos)	✓ (12–15 mos)	✓ (6–18 mos)	✓ (12–15 mos)		✓ (12–15 mos)	✓ (12–15 mos)	✓✓ (2 doses given 6 mos apart at age 12–23 mos)				
15 months													
18 months													
19–23 months	Catch-up ⁵	Catch-up ⁵	Catch-up ⁵ (to 5 years)	Catch-up ⁵	Catch-up ⁵ (to 5 years)		Catch-up ⁵	Catch-up ⁵				✓ ³ (given each fall or winter to children ages 6 mos–18 yrs)	
4–6 years		✓		✓			✓	✓					
7–10 years		Catch-up ⁵							Catch-up ⁵				
11–12 years		✓ Tdap			Catch-up ⁵			Catch-up ⁵	Catch-up ⁵		✓✓✓ ⁶		✓
13–18 years		Catch-up ⁵ (Tdap/Td)								Catch-up ^{5,6}	Catch-up ^{5,7}		

1. Your infant may not need a dose of HepB at age 4 months depending on the type of vaccine that your healthcare provider uses.
2. Your infant may not need a dose of Hib vaccine or RV vaccine at age 6 months depending on the type of vaccine that your healthcare provider uses.
3. One dose is recommended for most people. Children younger than age 9 years who are receiving influenza vaccine for the first time, or who received only 1 dose in the previous season (if it was their first vaccination season), should receive 2 doses spaced at least 4 weeks apart this season.
4. This dose of DTaP may be given as early as age 12 months if it has been 6 months since the previous dose.

5. If your child's vaccinations are delayed or missed entirely, they should be given as soon as possible.
6. All girls and women age 11 through 26 years should be vaccinated with 3 doses of HPV vaccine, given over a 6-month period. Boys and men age 11 through 26 years may also be vaccinated with one of the HPV vaccines (Gardasil) to reduce their likelihood of getting genital warts. The vaccine may be given to children as young as age 9 years.
7. If you have a teenager who is enrolling in college and planning to live in a dormitory and who hasn't previously been vaccinated against meningococcal disease, they should be vaccinated now.

Please note: Some children may need additional vaccines. Talk to your healthcare provider.

Recommended Immunization Schedule for Persons Aged 0 Through 6 Years—United States • 2010

For those who fall behind or start late, see the catch-up schedule

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years
Hepatitis B ¹		HepB	HepB				HepB					
Rotavirus ²				RV	RV	RV ²						
Diphtheria, Tetanus, Pertussis ³				DTaP	DTaP	DTaP	^{see footnote³}	DTaP				DTaP
<i>Haemophilus influenzae</i> type b ⁴				Hib	Hib	Hib ⁴	Hib					
Pneumococcal ⁵				PCV	PCV	PCV	PCV				PPSV	
Inactivated Poliovirus ⁶				IPV	IPV		IPV					IPV
Influenza ⁷							Influenza (Yearly)					
Measles, Mumps, Rubella ⁸							MMR		^{see footnote⁸}			MMR
Varicella ⁹							Varicella		^{see footnote⁹}			Varicella
Hepatitis A ¹⁰							HepA (2 doses)				HepA Series	
Meningococcal ¹¹											MCV	

Range of recommended ages for all children except certain high-risk groups

Range of recommended ages for certain high-risk groups

This schedule includes recommendations in effect as of December 15, 2009. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Considerations should include provider assessment, patient preference, and the potential for adverse

events. Providers should consult the relevant Advisory Committee on Immunization Practices statement for detailed recommendations: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS) at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Hepatitis B vaccine (HepB). (Minimum age: birth)

At birth:

- Administer monovalent HepB to all newborns before hospital discharge.
- If mother is hepatitis B surface antigen (HBsAg)-positive, administer HepB and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth.
- If mother's HBsAg status is unknown, administer HepB within 12 hours of birth. Determine mother's HBsAg status as soon as possible and, if HBsAg-positive, administer HBIG (no later than age 1 week).

After the birth dose:

- The HepB series should be completed with either monovalent HepB or a combination vaccine containing HepB. The second dose should be administered at age 1 or 2 months. Monovalent HepB vaccine should be used for doses administered before age 6 weeks. The final dose should be administered no earlier than age 24 weeks.
 - Infants born to HBsAg-positive mothers should be tested for HBsAg and antibody to HBsAg 1 to 2 months after completion of at least 3 doses of the HepB series, at age 9 through 18 months (generally at the next well-child visit).
 - Administration of 4 doses of HepB to infants is permissible when a combination vaccine containing HepB is administered after the birth dose. The fourth dose should be administered no earlier than age 24 weeks.
- ## 2. Rotavirus vaccine (RV). (Minimum age: 6 weeks)
- Administer the first dose at age 6 through 14 weeks (maximum age: 14 weeks 6 days). Vaccination should not be initiated for infants aged 15 weeks 0 days or older.
 - The maximum age for the final dose in the series is 8 months 0 days
 - If Rotarix is administered at ages 2 and 4 months, a dose at 6 months is not indicated.
- ## 3. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP). (Minimum age: 6 weeks)
- The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.
 - Administer the final dose in the series at age 4 through 6 years.
- ## 4. *Haemophilus influenzae* type b conjugate vaccine (Hib). (Minimum age: 6 weeks)
- If PRP-OMP (PedvaxHIB or Comvax [HepB-Hib]) is administered at ages 2 and 4 months, a dose at age 6 months is not indicated.
 - TriHibit (DTaP/Hib) and Hiberix (PRP-T) should not be used for doses at ages 2, 4, or 6 months for the primary series but can be used as the final dose in children aged 12 months through 4 years. See *MMWR* 1997;46(No. RR-8).
- ## 5. Pneumococcal vaccine. (Minimum age: 6 weeks for pneumococcal conjugate vaccine [PCV]; 2 years for pneumococcal polysaccharide vaccine [PPSV])
- PCV is recommended for all children aged younger than 5 years. Administer 1 dose of PCV to all healthy children aged 24 through 59 months who are not completely vaccinated for their age.
 - Administer PPSV 2 or more months after last dose of PCV to children aged 2 years or older with certain underlying medical conditions, including a cochlear implant.

6. Inactivated poliovirus vaccine (IPV). (Minimum age: 6 weeks)

- The final dose in the series should be administered on or after the fourth birthday and at least 6 months following the previous dose.
- If 4 doses are administered prior to age 4 years a fifth dose should be administered at age 4 through 6 years. See *MMWR* 2009;58(30):829–30.

7. Influenza vaccine (seasonal). (Minimum age: 6 months for trivalent inactivated influenza vaccine [TIV]; 2 years for live, attenuated influenza vaccine [LAIV])

- Administer annually to children aged 6 months through 18 years.
- For healthy children aged 2 through 6 years (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or TIV may be used, except LAIV should not be given to children aged 2 through 4 years who have had wheezing in the past 12 months.
- Children receiving TIV should receive 0.25 mL if aged 6 through 35 months or 0.5 mL if aged 3 years or older.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.
- For recommendations for use of influenza A (H1N1) 2009 monovalent vaccine see *MMWR* 2009;58(No. RR-10).

8. Measles, mumps, and rubella vaccine (MMR). (Minimum age: 12 months)

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.

9. Varicella vaccine. (Minimum age: 12 months)

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.
- For children aged 12 months through 12 years the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.

10. Hepatitis A vaccine (HepA). (Minimum age: 12 months)

- Administer to all children aged 1 year (i.e., aged 12 through 23 months). Administer 2 doses at least 6 months apart.
- Children not fully vaccinated by age 2 years can be vaccinated at subsequent visits
- HepA also is recommended for older children who live in areas where vaccination programs target older children, who are at increased risk for infection, or for whom immunity against hepatitis A is desired.

11. Meningococcal vaccine. (Minimum age: 2 years for meningococcal conjugate vaccine [MCV4] and for meningococcal polysaccharide vaccine [MPSV4])

- Administer MCV4 to children aged 2 through 10 years with persistent complement component deficiency, anatomic or functional asplenia, and certain other conditions placing them at high risk.
- Administer MCV4 to children previously vaccinated with MCV4 or MPSV4 after 3 years if first dose administered at age 2 through 6 years. See *MMWR* 2009; 58:1042–3.

Recommended Immunization Schedule for Persons Aged 7 Through 18 Years—United States • 2010

For those who fall behind or start late, see the schedule below and the catch-up schedule

Vaccine ▼	Age ►	7–10 years	11–12 years	13–18 years
Tetanus, Diphtheria, Pertussis ¹			Tdap	Tdap
Human Papillomavirus ²		see footnote 2	HPV (3 doses)	HPV series
Meningococcal ³		MCV	MCV	MCV
Influenza ⁴			Influenza (Yearly)	
Pneumococcal ⁵			PPSV	
Hepatitis A ⁶			HepA Series	
Hepatitis B ⁷			HepB Series	
Inactivated Poliovirus ⁸			IPV Series	
Measles, Mumps, Rubella ⁹			MMP Series	
Varicella ¹⁰			Varicella Series	

Range of recommended ages for all children except certain high-risk groups

Range of recommended ages for catch-up immunization

Range of recommended ages for certain high-risk groups

This schedule includes recommendations in effect as of December 15, 2009. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Considerations should include provider assessment, patient preference, and the potential for adverse

events. Providers should consult the relevant Advisory Committee on Immunization Practices statement for detailed recommendations: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS) at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap). (Minimum age: 10 years for Boostrix and 11 years for Adacel)

- Administer at age 11 or 12 years for those who have completed the recommended childhood DTP/DTaP vaccination series and have not received a tetanus and diphtheria toxoid (Td) booster dose.
- Persons aged 13 through 18 years who have not received Tdap should receive a dose.
- A 5-year interval from the last Td dose is encouraged when Tdap is used as a booster dose; however, a shorter interval may be used if pertussis immunity is needed.

2. Human papillomavirus vaccine (HPV). (Minimum age: 9 years)

- Two HPV vaccines are licensed: a quadrivalent vaccine (HPV4) for the prevention of cervical, vaginal and vulvar cancers (in females) and genital warts (in females and males), and a bivalent vaccine (HPV2) for the prevention of cervical cancers in females.
- HPV vaccines are most effective for both males and females when given before exposure to HPV through sexual contact.
- HPV4 or HPV2 is recommended for the prevention of cervical precancers and cancers in females.
- HPV4 is recommended for the prevention of cervical, vaginal and vulvar precancers and cancers and genital warts in females.
- Administer the first dose to females at age 11 or 12 years.
- Administer the second dose 1 to 2 months after the first dose and the third dose 6 months after the first dose (at least 24 weeks after the first dose).
- Administer the series to females at age 13 through 18 years if not previously vaccinated.
- HPV4 may be administered in a 3-dose series to males aged 9 through 18 years to reduce their likelihood of acquiring genital warts.

3. Meningococcal conjugate vaccine (MCV4).

- Administer at age 11 or 12 years, or at age 13 through 18 years if not previously vaccinated.
- Administer to previously unvaccinated college freshmen living in a dormitory.
- Administer MCV4 to children aged 2 through 10 years with persistent complement component deficiency, anatomic or functional asplenia, or certain other conditions placing them at high risk.
- Administer to children previously vaccinated with MCV4 or MPSV4 who remain at increased risk after 3 years (if first dose administered at age 2 through 6 years) or after 5 years (if first dose administered at age 7 years or older). Persons whose only risk factor is living in on-campus housing are not recommended to receive an additional dose. See *MMWR* 2009;58:1042–3.

4. Influenza vaccine (seasonal).

- Administer annually to children aged 6 months through 18 years.
- For healthy nonpregnant persons aged 7 through 18 years (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or TIV may be used.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.
- For recommendations for use of influenza A (H1N1) 2009 monovalent vaccine. See *MMWR* 2009;58(No. RR-10).

5. Pneumococcal polysaccharide vaccine (PPSV).

- Administer to children with certain underlying medical conditions, including a cochlear implant. A single revaccination should be administered after 5 years to children with functional or anatomic asplenia or an immunocompromising condition. See *MMWR* 1997;46(No. RR-8).

6. Hepatitis A vaccine (HepA).

- Administer 2 doses at least 6 months apart.
- HepA is recommended for children older than 23 months of age who live in areas where vaccination programs target older children or who are at increased risk for infection or for whom immunity against hepatitis A is desired.

7. Hepatitis B vaccine (HepB).

- Administer the 3-dose series to those not previously vaccinated.
- A 2-dose series (separated by at least 4 months) of adult formulation Recombivax HB is licensed for children aged 11 through 15 years.

8. Inactivated poliovirus vaccine (IPV).

- The final dose in the series should be administered on or after the fourth birthday and at least 6 months following the previous dose.
- If both OPV and IPV were administered as part of a series, a total of 4 doses should be administered, regardless of the child's current age.

9. Measles, mumps, and rubella vaccine (MMR).

- If not previously vaccinated, administer 2 doses or the second dose for those who have received only 1 dose, with at least 28 days between doses.

10. Varicella vaccine.

- For persons aged 7 through 18 years without evidence of immunity (see *MMWR* 2007;56[No. RR-4]), administer 2 doses if not previously vaccinated or the second dose if only 1 dose has been administered.
- For persons aged 7 through 12 years, the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.
- For persons aged 13 years and older, the minimum interval between doses is 28 days.